

Food Allergy Action Plan Step 1: Prevention

Camper's Name: _____ Date of Birth: _____
Teacher's Name: _____ Room #: _____
ALLERGY TO: _____
Asthmatic? Y/N) _____ (Yes=Higher Risk for Severe Reaction)

Camp will:

- ☐ A Certified Medication Technician on site with on-call Delegating RN
- ☐ Have staff trained in CPR & First Aid
- ☐ Have staff trained in Allergy & Anaphylaxis
→ administering EpiPen® including demonstration & practice
- ☐ Emergency List distributed to: _____
- ☐ Have staff trained on individual emergency plans
- ☐ School staff will make every reasonable effort to prevent the student's exposure to known allergens
- ☐ Other _____

Parents will:

- ☐ Provide pertinent health information to the camp
- ☐ Provide Physician Authorization Forms and Action Plans
→ for student medication and specific actions plans for emergency care
- ☐ Current, non-expired medications
- ☐ Provide safe snack option to camp
- ☐ Other: _____
- ☐ Other: _____
- ☐ Other: _____

Student will:

- ☐ Make every effort to avoid contact with allergen
- ☐ Alert nearest adult if suspect exposure to allergen
- ☐ Other _____

Notes:

Food Allergy Action Plan



Camper's Name: _____

Date of Birth: _____

Teacher's Name: _____

Room #: _____

ALLERGY TO: _____

Asthmatic? (Y/N) _____ (Yes=Higher Risk for Severe Reaction)

STEP 2: TREATMENT

Symptoms	Give This Medication	
	Epinephrine	Antihistamine
If a food allergen is ingested or suspected bee sting, but <i>no symptoms</i>		
Mouth: itching, tingling, or swelling of lips, tongue mouth		
Skin: hives, itchy rash, swelling of the face or extremities		
Gut: nausea, abdominal cramps, vomiting, diarrhea		
Throat *: Tightening of throat, hoarseness, hacking cough		
Lung*: Shortness of breath, repetitive coughing, wheezing		
Heart*: Weak or thread pulse, low blood pressure, fainting, pale, blueness		
Other:		
If reaction is progression (several of the above areas affected):		

*Potentially life-threatening. The severity of symptoms can quickly change.

Administer:

_____ Epinephrine auto-injector 0.15 mg

_____ Epinephrine auto-injector 0.3 mg

_____ Repeat dose if EMS has not arrived in 10 minutes

Antihistamine: give _____

Other: give _____

IMPORTANT: CALL 911 IMMEDIATELY
Asthma inhalers and/or antihistamines can not be depended on to treat anaphylaxis.

STEP 3: EMERGENCY CONTACTS

_____	_____
Parent's Name	Phone Number
_____	_____
Secondary Emergency Contact	Phone Number
_____	_____
Consulting School Nurse	Phone Number

Parent Guardian's Signature/Date

Doctor's Signature/Date